

Stefanie Mikulics, MD  
1050 Las Tablas Road, Suite 2  
Templeton, CA 93465  
805-434-9441

#### DERMAL FILLERS: PRE TREATMENT INSTRUCTIONS

\* 3 DAYS BEFORE treatment: AVOID topical products such as Tretinoin (Retin-A), Retinols, Retinoids, Glycolic Acid, Alpha Hydroxy Acid, or any “anti-aging” products. Also AVOID waxing, bleaching, tweezing, or the use of hair removal cream on the area to be treated.

NO DENTAL WORK can be done 30 days BEFORE or AFTER any Filler Treatments. Please note: If you have a cardiovascular history, please check with your doctor prior to stopping use of Aspirin.

\* 7 DAYS BEFORE treatment (to prevent bruising): AVOID blood thinning over-the-counter medications such as Aspirin, Motrin, Ibuprofen, and Aleve. Also avoid herbal supplements, such as Garlic, Vitamin E, Ginkgo Biloba, St. John’s Wort, and Omega-3 capsules.

Please note: If you have a cardiovascular history, please check with your doctor prior to stopping use of Aspirin.

\* Do not drink alcoholic beverages 24 hours before (or after) your treatment to avoid extra bruising.

Inform your provider if you have a history of Perioral Herpes to receive advice on antiviral therapy prior to treatment.

\* Do not use Dermal Fillers if you are pregnant or breastfeeding, are allergic to any of its ingredients, or suffer from any neurological disorders. Please inform your provider if you have any questions about this prior to the treatment.

#### **Day of Treatment**

\* Arrive to the office with a “clean face”. Please do not wear makeup. You may bring your own makeup to apply after your treatment.

\* To maximize your comfort during the procedure, a topical anesthetic may be applied. In some cases, a local numbing medication will be injected into or around the area(s) to be treated.

\* You may experience a mild amount of tenderness or a stinging sensation following injection.

\* To ensure a smooth and even correction, your provider may massage the area(s) treated, which may cause a temporary, minimal amount of redness to your skin.

### **Immediately After Treatment**

\* Redness and swelling are normal. Bruising may also be visible.

\* You may experience some tenderness at the treatment site(s) that can last for a few hours or a few days.

\* Depending upon the area(s) treated and product(s) used, you may feel “firmness” in the treated area(s) for 1 to 2 weeks after treatment. Over time, the area(s) will soften and “settle”.

### **DERMAL FILLERS: POST TREATMENT INSTRUCTIONS**

\* Apply an ice or cold gel pack to the area(s) treated (avoiding pressure) as this helps reduce swelling and the potential for bruising.

\* Once you have adequately cooled/iced the area(s) as instructed and any pinpoint bleeding from the injection site(s) has subsided, you may begin wearing makeup.

\* AVOID placing excessive pressure on the treated area(s) for the first few hours and up to 2 to 3 days; when cleansing your face or applying makeup, be very gentle.

\* AVOID exercise or strenuous activities for the remainder of the treatment day; you may resume other normal activities/routines immediately.

\* TRY sleeping on your back for the first few nights after treatment to avoid possible manipulation to the treated area. (Putting a pillow underneath your knees can help keep you on your back.)

\* You may take Acetaminophen/Tylenol if you experience any mild tenderness or discomfort.

NO DENTAL WORK can be done 30 days BEFORE or AFTER any Filler Treatments. Please note: If you have a cardiovascular history, please check with your doctor prior to stopping use of Aspirin.

\* AVOID topical products such as Tretinoin (Retin-A), Retinols, Retinoids, Glycolic Acid, Alpha Hydroxy Acid, or any “anti-aging” products. Also AVOID waxing, bleaching, tweezing, or the use of hair removal cream on the area to be treated for two (2) days after treatment. It is best to wear no make-up or lipstick until the next day. Earlier can cause pustules.

\* AVOID extended UV exposure until any redness/swelling has subsided. Be sure to apply an SPF 30 or higher sunscreen.

\* Wait a minimum of four weeks (or as directed by your provider) before receiving any skin care or laser treatments.

\* Untreated bruising will generally fade in 5 to 14 days.

\* Maximum results usually take 2 full weeks after initial treatment; therefore, make no judgement about your results until the full two weeks have occurred. You must wait until then before retreating or correcting.

\*\*\*\*\*Please report any redness, blisters, or itching immediately if it occurs after treatment.

I certify that I have been counseled in post treatment instructions and have been given written instructions as well.

---

Signature

Date